



**Evidence-Based Clinical Assessment Toolkit  
(EBCAT)**

**Behavioural Expressions  
Workbook**



# COPYRIGHT

© 2014 Applied Aged Care Solutions Pty Ltd ABN 46 083 264 359

All rights reserved. Applied Aged Care Solutions Pty Ltd (**AACS**):

- ❖ invests significant time and resources in creating its original products;
- ❖ protects its copyright material;
- ❖ will enforce its rights in copyright material;
- ❖ reserves its legal rights to claim its loss and damage or an account of profits made resulting from infringements of its copyright.

Except as otherwise set out, AACS owns all copyright in this pack (“product”). AACS grants **Barwon Health** and the **Department of Health Victoria** a non-exclusive, perpetual, irrevocable, non-transferable, worldwide and royalty free licence to use the materials.

Except as permitted by the Copyright Act 1968 (Cth) or unless you have obtained the specific written permission of AACS, Barwon Health, or the Department of Health Victoria, **you must not**, whether by electronic, mechanical or graphical means, or by hand or by any other means:

- reproduce or copy this product in whole or in part;
- modify or adapt this product;
- use this product for training purpose without licence;
- publish this product in whole or in part;
- cause this product in whole or in part to be transmitted;
- store this product in whole or in part in a retrieval system including a computer;
- record this product in whole or in part either electronically or mechanically; or
- resell this product in whole or in part.

# DISCLAIMER

Applied Aged Care Solutions Pty Ltd (**AACS**) and its authors are not responsible for the results of any action taken on the basis of any information in this publication or for any error in or omission from this publication.

This publication contains information of a general nature that may not be suitable to your own circumstances. It is your responsibility to satisfy yourself as to its suitability for your circumstances.

The publication has been compiled from information provided by third parties and other sources. There may be technical inaccuracies, typographical errors or unsuitable material for you. If you have any concerns about such third party content, you should refer to those sources for more specialist information and advice. Attribution to such sources appears in the text of this publication.

To the fullest extent permitted by law, AACS and its authors will be under no liability to any person in respect of any loss or damage (including consequential loss or damage) which may be suffered or incurred or which may arise directly or indirectly from or in connection with any use of this publication or the information contained in it.

To the fullest extent permitted by law, all express or implied warranties are hereby excluded. Warranties or guarantees implied by statute which cannot be excluded will apply, however the liability of AACS and its authors for a breach of such implied terms will be limited, at the option of AACS, to any one or more of the following:

- replacement of the product supplied or supply of an equivalent product;
- payment of the cost of replacing the product or of acquiring an equivalent product;
- payment of the cost of having the product rectified;
- and in no circumstances extends to consequential, indirect or special loss or damage.

# Acknowledgements

Barwon Health was funded by the Department of Health Victoria to develop an evidence based Aged Care Funding Instrument (ACFI) toolkit and associated process guidelines for public sector residential aged care services (PSRACS). Barwon Health engaged Applied Aged Care Solutions (AACS), developers of the original ACFI to conduct the project and develop the toolkit. AACS has extensive experience in designing assessment models and evidence-based tools for residential aged care.

Additional content matter expertise was provided to assist with the development of the materials. Applied Aged Care Solutions (AACS) worked with:

## **La Trobe University**

Dr Deirdre Fetherstonhaugh (Director, Senior Research Fellow), Dr Michael Bauer (Senior Research Fellow ) and Dr. Margaret Winbolt (Senior Research Fellow, Director Victoria and Tasmania Dementia Training Study Centre ) from the Australian Centre for Evidence Based Aged Care (ACEBAC), Australian Institute for Primary Care and Ageing (AIPCA), La Trobe University provided feedback on the EBCAT framework and the process content of the workbooks (ADL, Continence, Medicines and Complex Health Workbooks).

## **Monash University**

Professor Daniel O'Connor (Professor of Old Age Psychiatry, Faculty of Medicine, Nursing & Health Sciences, and Head of the Aged Mental Health Research Unit Monash University) provided feedback on the framework and the process content of the Cognitive and Behavioural Expressions Workbooks.

## **PRACS Leadership Group**

AACS acknowledges the comments of the PSRACS Nurse Leadership Group and feedback and comment provided at the workshops for PSRACS staff hosted by the Department of Health Victoria.

## **Other Contributors**

Professor Rhonda Nay who reviewed the Complex Health Workbook.

# Contents

Acknowledgements.....	3
List of Tables .....	5
List of Figures .....	5
Acronyms and Glossary.....	6
Overview of the Toolkit Products .....	8
Suggested Roles for Staff Implementing the Toolkit .....	9
Introduction to the Behavioural Expressions Workbook .....	10
Topic 6:.....	12
Behavioural Expressions .....	12
The Behavioural Expressions Topic.....	13
Investigating Behavioural Expressions.....	13
Getting Started with a Behavioural Example.....	17
Preparation .....	18
Recommended Staffing Skill Set .....	18
Background Reading .....	19
Some Basics.....	21
Behavioural Symptoms .....	23
Psychological Symptoms.....	24
Psychiatric Symptoms .....	25
How Behavioural Expressions Interact with Other Domains.....	27
Identification Process.....	29
Gathering the History .....	29
Documentation .....	29
Resident and Family .....	31
Screening.....	31
Comprehensive Assessment .....	35
For wandering, verbal or physical behaviour .....	37
For a mood symptom.....	40
Clinical Risks .....	41
Implementation Phase.....	43
Referrals.....	43

Interventions.....	44
Non Pharmacological Approaches.....	45
Communicating Effectively.....	46
Care Planning.....	49
Goal Setting.....	49
Role of Documentation.....	49
Linking the Evidence.....	51
Completing the ACFI documentation.....	52
Evaluation Process.....	54
Outcomes.....	54
Summary: Steps and Information Flow.....	55
Behavioural Expressions Resources.....	57
Behavioural Expressions References.....	58
Behavioural Expressions Workbook Exercises.....	60

## List of Tables

Table 1: Staff Activities for the Behavioural Expressions Process.....	18
Table 2: Modified extract from the Behavioural Descriptions List.....	33
Table 3: Referral for Behavioural Expressions.....	44
Table 4: ACFI Questions 7 to 9 Checklists.....	52
Table 5: ACFI Question 10 Checklist.....	53
Table 6: Behavioural Expressions Workbook Resources.....	57
Table 7: References for the Behavioural Expressions Workbook.....	58

## List of Figures

Figure 1: Behavioural Expressions Process.....	16
Figure 2: Behavioural Assessment Process.....	34
Figure 3: Behavioural Expressions Summary: Steps and information flow.....	56

## Acronyms and Glossary

<b>ACCR</b>	Aged Care Client Record
<b>ACFI</b>	Aged Care Funding Instrument
<b>ADL</b>	Activities of Daily Living
<b>BAF</b>	Behavioural Assessment Form
<b>BCOPE</b>	Best Care For Older People Everywhere
<b>BPSD</b>	Behavioural and Psychological Symptoms of Dementia
<b>CAM</b>	Confusion Assessment Method
<b>CCF</b>	Care Continuum Framework
<b>CDAMS</b>	Cognitive, Dementia and Memory Service
<b>CDC</b>	Consumer Directed Care
<b>CHA</b>	Comprehensive Health Assessment (CHA) for Older People in the Health Care System
<b>CHAOP</b>	Comprehensive Health Assessment of the Older Person
<b>DBMAS</b>	Dementia Behaviour Management Advisory Service
<b>DOMS</b>	Dementia Outcomes Measurement Suite
<b>EBCAT</b>	Evidence Based Clinical Assessment Toolkit
<b>EBCAT Assessment Packs</b>	Each workbook has an assessment pack. This contains the recommended screens, assessments and Standardised Care Processes recommended within the workbook.
<b>EBCAT Introductory Guide</b>	This document presents: Project methodology; Overview of products; and details of the Management role;
<b>EBCAT Reading Pack</b>	This document provides the background reading for all EBCAT Workbooks.
<b>EBCAT Topics</b>	<ol style="list-style-type: none"> <li>1. Nutrition</li> <li>2. Mobility</li> <li>3. Self-care (Personal Hygiene, Toileting)</li> <li>4. Continence</li> <li>5. Cognition</li> <li>6. Behavioural Expressions (Wandering, Verbal &amp; Physical, Mood)</li> <li>7. Medicines</li> <li>8. Pain</li> <li>9. Swallowing</li> <li>10. Skin &amp; Wounds</li> </ol>
<b>EBCAT Workbooks</b>	<p>The toolkit is presented in six ‘user friendly educational workbooks’ to walk the user through the process of using evidence-based clinical assessment tools for each domain of:</p> <ul style="list-style-type: none"> <li>• ADL Workbook (Topics 1-3)</li> <li>• Continence Workbook (Topic 4)</li> <li>• Cognition Workbook (Topic 5)</li> <li>• Behavioural Expressions Workbook (Topic 6)</li> <li>• Medicine Workbook (Topic 7)</li> </ul>

	<ul style="list-style-type: none"> <li>• Complex Health Workbook (Topics 8-10)</li> </ul>
<b>FRAT</b>	Falls Risk Assessment Tool
<b>GP</b>	General Practitioner
<b>IPA</b>	International Psychogeriatric Association
<b>KICA-Cog</b>	Kimberley Indigenous Cognitive Assessment
<b>MP</b>	Medical Practitioner
<b>M-VRBPI</b>	Modified Resident Verbal Brief Pain Inventory
<b>NATFRAME</b>	National Framework for Documenting Care in Residential Aged Care Services <a href="http://webarchive.nla.gov.au/gov/20140803082152/http://www.health.gov.au/internet/publications/publishing.nsf/Content/ageing-rescare-natframe.htm~ageing-rescare-natframe01.htm">http://webarchive.nla.gov.au/gov/20140803082152/http://www.health.gov.au/internet/publications/publishing.nsf/Content/ageing-rescare-natframe.htm~ageing-rescare-natframe01.htm</a>
<b>NCD</b>	Neuro-Cognitive Disorder
<b>NPI-NH</b>	Neuro-Psychiatric Inventory for Nursing Homes
<b>NRS</b>	Numeric Pain Rating Scale
<b>PAINAID</b>	Pain Assessment in Advanced Dementia
<b>PAS-CIS</b>	Psychogeriatric Assessment Scales- Cognitive Impairment Scale
<b>PCC</b>	Person Centred Care
<b>PMS</b>	Physical Mobility Scale
<b>PSRACS</b>	Public Sector Residential Aged Care Services
<b>QoC</b>	Quality of Care
<b>QoL</b>	Quality of Life
<b>RACF</b>	Residential Aged Care Facilities
<b>RNDC</b>	Resident Nutrition Data Card
<b>ROM</b>	Range Of Movement
<b>RUDAS</b>	Rowland Universal Dementia Assessment Scale
<b>SCP</b>	Standardised Care Process
<b>SCORE</b>	Strengthening Care Outcomes for Residents with Evidence
<b>VDS</b>	Verbal Descriptor Scale

## Overview of the Toolkit Products

The Evidence-Based Clinical Assessment Toolkit (EBCAT) consists of the following products:

Resource	How used
Introductory Guide	<p>The Introductory Guide is aimed at the lead nurse and the nursing management.</p> <p>It presents: the project methodology; an introduction to the products; and details on the Nursing Management role.</p>
Reading Pack	<p>The Reading Pack, provides further reference information for the background reading section of each workbook, it is aimed at care staff.</p> <p>This pack contains reading material which cannot be sourced from the internet. References for supporting material that can be sourced off the internet are provided in workbook appendices. There is also a sample Quality of Life questionnaire in the Reading Pack.</p>
Workbooks	<p>The EBCAT Workbooks are designed to be used by the lead nurse. The workbooks should be used as a training tool by the lead nurse when training the care staff on the EBCAT. There are six workbooks which cover the domains of:</p> <ul style="list-style-type: none"> <li>• Activities of Daily Living</li> <li>• Continence</li> <li>• Cognition</li> <li>• Behavioural Expressions</li> <li>• Medicine</li> <li>• Complex Health</li> </ul> <p>Each workbook contains detailed information and case studies on how to complete the recommended assessment tools as part of a nursing-based process. The Appendices provide references for the suggested resources, and a workbook exercise to practice what has been learnt.</p>
Quick Guides	<p>The Quick Guides are designed for use by care staff. There is one quick guide per workbook.</p> <p>The Quick Guide is a quick reference to the EBCAT process and tools. It is recommended it be kept handy for use on the 'floor', whenever required.</p>
Assessment Packs	<p>The assessment packs contains the recommended screen, assessment tools and relevant clinical risk tools. There is one assessment pack per workbook.</p> <p>The tools are used by the care staff when identifying the needs of the residents.</p>



## Suggested Roles for Staff Implementing the Toolkit

The toolkit requires the participation of three types of staff as described below.

	<b>Who</b> and what they do in regard to the Toolkit
<b>Nursing Management</b>	<p>This group would typically consist of nursing staff who do not work ‘on the floor’, for example the Director of Nursing or Nurse Unit Manager. They are vital to ensuring, that the toolkit is set up properly to support implementation at the site, to ensure that the process is continuously monitored and improved, and to monitor the process to ensure the documentation and ACFI claiming is accurate.</p> <p>The nursing management role includes:</p> <ul style="list-style-type: none"> <li>○ Preparing the toolkit and auditing for readiness to implement</li> <li>○ Selecting a lead nurse for the leadership role and to train the care staff</li> <li>○ Implementing the toolkit and monitoring the progress</li> </ul> <p>The nursing management role is described in detail in the Introductory Guide.</p>
<b>Lead Nurse</b>	<p>This person will be selected by the Nursing Management group to lead the EBCAT process at the site. It is recommended they be a nurse (RN or EN).</p> <p>The lead nurse role includes:</p> <ul style="list-style-type: none"> <li>○ Assisting the Nursing Management group to prepare the toolkit</li> <li>○ Training the care staff on how to implement the EBCAT process and tools</li> <li>○ Providing leadership to the care staff during the implementation of the process</li> <li>○ Assisting the Nursing Management group to monitor the progress</li> </ul> <p>The lead nurse role is described in detail in the Introductory Guide.</p>
<b>Care Staff</b>	<p>This group are the nurses (RN or EN) and Personal Care Workers who deliver the daily care to the residents ‘on the floor’.</p> <p>They receive the training and implement the EBCAT process and tools when undertaking the resident assessment process.</p>

## Introduction to the Behavioural Expressions Workbook

The Behavioural Expressions Workbook is one of six that form the Evidence Based Clinical Assessment Toolkit (EBCAT). This workbook is one of four resources relevant to the Behavioural Expressions domain which comprise:

- A Reading Pack
- Behavioural Expressions Workbook
- Behavioural Expressions Quick Guide
- Behavioural Expressions Assessment Pack

The toolkit aims to provide a resource to assist Public Sector Residential Aged Care Services (PSRACS) staff to systematically and consistently determine and manage resident care needs. The toolkit uses evidence-based clinical assessment tools for assessing and managing residents with the goals of improving the clinical and quality of life for the residents and demonstrating accountability to government regulators for example, with the Aged Care Funding Instrument (ACFI) requirements.

During 2013, the Australian Government made changes to the Aged Care Funding Instrument (ACFI) requiring further evidence to support funding claims made by services with activities of daily living support needs. In addition, the Australian government introduced more stringent penalties for providers with inaccurate or misleading ACFI appraisals from 1 July 2013.

This workbook will assist your service to meet the ACFI evidence requirements for the three ACFI behaviours items (ACFI 7- Wandering; ACFI 8- Verbal Behaviour; ACFI 9- Physical Behaviour) and the depression item (ACFI 10). The workbook uses familiar and freely available Australian toolkits and resources including:

- ACFI Assessment pack (Behaviour Records, Cornell Scale for Depression in Dementia)
- ACFI User Guide
- Neuropsychiatric Inventory- Nursing Home Version (NPI-NH)
- An Initial Nurse Assessment – e.g., the Comprehensive Health Assessment (CHA) for Older People in the Health Care System, which was designed for recording assessment results based on the Comprehensive Health Assessment of the Older Person (CHAOP) resource
- The NATFRAME (National Framework for Documenting Care in Residential Aged Care Services)

## Behavioural Expressions Topic

- The International Psychogeriatric Association (IPA) Guides to Behavioural and Psychological Symptoms of Dementia (BPSD)
- Dementia Outcomes Measurement Study (DOMS)
- Best Care for Older People Everywhere (BCOPE). The toolkit.
- Standardised Care Processes (SCP Physical Restraint and SCP Depression)
- Diagnostic and Statistical Manual of Mental Disorders fifth edition (DSM-5)

# **Topic 6:**

# **Behavioural Expressions**

## The Behavioural Expressions Topic

This topic covers behavioural symptoms associated with cognitive impairment disorders as well as psychological/neuro-psychiatric symptoms (e.g. depressive symptoms, hallucinations, delusions etc).

Behaviour is the externally visible activity, response or expression of a person to an internal and/or external stimuli or unmet need (i.e., something they perceive or is actually happening or a form of communication related to an unmet need). The activity, response or behavioural expression is determined by the way a person interprets and processes the stimuli or communicates the unmet need. A person with moderate or severe cognitive impairment or dementia will have a limited ability to make a rationale or logical interpretation of the stimuli or how to communicate an unmet need. The behavioural expressions can be considered a form of communication (Kozman et al, 2006) but in the case of a person with dementia it will not form a volitional, deliberate or rationale course of actions designed to produce a particular outcome.

The characterisation of a behaviour as problematic in the case of residential aged care is contextual and framed by the prevailing societal norms, expectations, views and attitudes of others such as other residents, the family/visitors and the staff who are supporting the person on a daily basis.

When evaluating behavioural expressions it is important to understand the contributions of the:

- (i) social and physical environment
- (ii) care routines
- (iii) staff and resident relationships and interactions
- (iv) the needs of the person and how they try and communicate; and
- (v) person's broader health care needs

Any analysis of the reasons for problematic behavioural expressions requires a comprehensive understanding of the 'trigger' events to determine the contributing factors to the behavioural expression.

## Investigating Behavioural Expressions

The following four process steps should be followed when investigating behavioural expressions (consistent across all EBCAT topics). The steps are:

**1. Preparation of staff** – ensuring that staff have the required qualifications or competencies and have completed background reading if required. Staff will need to have a good level of background knowledge of possible psychological and neuro-cognitive behavioural symptoms, what they are and how they impact on people. Resources incorporated into the background reading section include:

- The International Psychogeriatric Association Guidelines
- Best Care for Older People Everywhere (BCOPE). The Toolkit (2012)
- On-line education module “Evidence-based tools for BPSD Assessment.

The references for these resources can be found in the Behavioural Expressions Appendix.

**2. Identifying** – gathering the resident’s history by collating documents and talking with residents and their family, looking for signs and symptoms in the initial nurse assessment, completing a comprehensive assessment approach and assessing the scope of the challenge. It is recommended that all new residents have a comprehensive assessment of nutrition. A comprehensive approach will include:

File Notes Review:

- Aged Care Client Record (ACCR) – Parts 4 and 5 provide information on cognitive, behaviour and psychological aspects
- Comprehensive Medical Assessment – which (if available) may have sections on behavioural, psychological and neuro-psychiatric symptoms

Screen:

- Initial nurse assessment – e.g., the CHA records if lowered mood is apparent
- A screening step across all behavioural expressions that will cover the ACFI items 7 to 10
- Supported with a behaviour assessment approach based on the literature recommended ‘ABC’ theory (the modified Behavioural Assessment Form)

Further Assessment:

The following assessment tools and Standardised Care Processes (found in the Behavioural Expressions Assessment Pack) are recommended for assessing the person’s behavioural needs.

- For behaviours (ACFI 7, ACFI 8, ACFI 9) the mandatory assessments are Behaviour Records

## Behavioural Expressions Topic

- For depression (ACFI 10) the mandatory assessment is the Cornell Scale of Depression in Dementia

The SCORE Standardised Care Process for addressing the clinical risk of delirium is recommended.


**3. Implementing** – based on the information from the identification phase this covers making needed referrals, selecting and trying interventions, writing a care plan, completing the ACFI and documenting the process. The steps are:

- Undertaking referral options to complete gaps or seek specialist advice
- Planning evidence-based care strategies to assist the person to maintain or possibly improve their participation ability
- Listening to and setting goals with the consumer (resident and family) to hear their understanding and personalise the approach
- Understanding the role of communicating the care needs, goals and strategies through care plans and resident file notes to provide the documentation trail
- Completing ACFI documentation

**4. Evaluating** – monitoring and evaluating the effectiveness of the process, interventions and looking for ways to further improve the care outcomes for residents.

The overall process and associated activities are illustrated in Figure 1 which shows the phases (e.g. prepare, identify, implement, evaluate), the steps within each phase and the aim for each phase (e.g. have identified the care needs of the resident). Each topic of the Behavioural Expressions Workbook follows the same pattern. Consistent application of this process will assist your home provide systematic, accountable care and facilitate the best possible resident care outcomes.

Figure 1: Behavioural Expressions Process





## Getting Started with a Behavioural Expressions Example

This case study will be referred to as we discuss the topic of behavioural expressions.

*Mr. Grey is 80 years old, married with one son, a retired plumber, whose hobby was amateur theatre acting and singing. He has advanced dementia, is independently mobile and manages his personal hygiene tasks with a little bit of prompting and setting up by his wife. Mrs. Grey manages the household (finances, domestic chores, the garden and transport etc).*


*It is his wandering away from home and verbal combativeness that is worrying Mrs. Grey. She described him as a “man on a mission”- walking out of the house most mornings and heading up the street. When Mrs. Grey tried to stop him, he often seemed not to recognise her, and she found him quite threatening when he yelled at her to get out of his way. Usually he returned within a few minutes forgetting why he left the house, and a cup of tea settled him down.*

*Mr. Grey’s urgent outings had become more frequent, and he is more restless and more unresponsive to his wife’s conversation or directions. There were a few times when he had to be returned by the police as he was unable to find his way home.*

*He was recently assessed by the Aged Care-Assessment Service and approved for residential care, and a bed has become available at a dementia specific aged care home.*



## Preparation



There are two specific aspects to **preparing** staff for understanding and assisting residents with behavioural expressions. They are:

- 1) Ensuring that staff have the required qualifications or competencies; and
- 2) Completing the pre-reading if required

### Recommended Staffing Skill Set

Table 1 below provides a structure for management to identify which staff have the skills required to complete activities within the behavioural expressions process. The process includes:

- Identifying the required activities (examples provided in Table 1)
- Assessing staff competency to complete the activities
- Addressing identified gaps

This will assist nursing management to select and determine the roles of staff to ensure the process can be completed effectively. For example, if there is a gap found in the behavioural assessment and management activity, the facility could consider further training of current staff, or securing a nurse with the required clinical knowledge, or identifying a local Allied Health Professional (i.e. Psychologist) who could complete the assessment.

The introductory guide also provides further instructions for nursing management in preparation for implementing this toolkit.

**Table 1: Staff Activities for the Behavioural Expressions Process**

Activity	Responsible for sign off	Does the activity
Collating Documents		
Identifying needs from the collation documents		
Initial Nurse Assessment: e.g. CHA		
Screening/Assessment : Modified BAF		

Activity	Responsible for sign off	Does the activity
Behavioural Description List		
Charting- Behaviour Records		
Assessment: Modified Cornell Scale of Depression in Dementia		
Documenting into file notes		
Determine and action Referrals		
Develop Interventions and Strategies		
Write up the Care Plan		
Complete ACFI associated documents		
Complete Standards or audit documents		
Review the Care Plan		
Evaluate Resident Care outcomes		

## Background Reading

Staff should complete the background reading or be able to demonstrate that they have adequate experience and knowledge of behavioural and psychiatric issues in older people. It is expected that staff will have:

- A comprehensive understanding of the behavioural expressions topic and how it is impacted by other health areas
- An awareness of associated diagnoses, signs and symptoms
- Knowledge about and training on how to use the assessment tools
- Knowledge about evidence informed practices associated with the behavioural expressions topic to assist with the development and implementation of evidence-based care plans

The background reading relevant to the behavioural expressions topic is:

- **IPA Complete Guides to Behavioural and Psychological Symptoms of Dementia (BPSD) –Specialists Guide and Nurses Guide.** The content is designed to contribute

to the improved management and support of residents with dementia and BPSD and to reduce some of the stresses experienced by care staff. The principle of Person Centred Care approach is widely supported in these guides. Each module includes an in-depth discussion and analysis of its subject area and concludes with a reference and recommended reading list.

The IPA Complete Guides to BPSD – Specialists Guide provides extensive background reading covering eight modules. “BPSD is recognizable, understandable and treatable. The recognition and appropriate management of BPSD are important factors in improving our care of dementia patients and their caregivers and is central to the development of The IPA Complete Guides to BPSD.”(Preface p. iii.)

The Specialists Guide has eight modules:

- Module 1 – An introduction to BPSD
- Module 2 – Clinical issues
- Module 3 – Etiology
- Module 4 – Role of family caregivers
- Module 5 – Non-pharmacological treatments
- Module 6 – Pharmacological management
- Module 7 – Cross-cultural and transnational considerations
- Module 8 – Long-term care

The related Nurses Guide is in a shorter, summary format, and is designed to complement the Specialists Guide.

- **Australian Dementia Training Study Centres on-line education module “Evidence-based tools for BPSD Assessment”**

The Australian Dementia Training Study Centres (DTSCs) provide a broad range of dementia related educational services to meet the needs of the health and aged care sector. The Queensland DTSC has developed an on-line education module “Evidence-based tools for BPSD Assessment” which informs on three topics and describes symptoms and their status, discusses common assessment tools that are evidence based, and discusses interventions/care planning for:

- Agitation/Aggression
- Apathy
- Wandering

References for these resources are found in the Behavioural Expressions Appendix.

## Some Basics

**Person-centred care** advocates the importance of language to change perceptions of people living in care. For example:

- It is not a “demented person,” “dementia patient” but “a person with dementia,” putting the person first
- It is not “suffering” from dementia; it is a person “living with dementia”
- Instead of describing actions of people with dementia for example as “stealing” which implies that these are deliberate actions we should use words which simply describe what was observed to happen without incorrectly labeling the person’s intent. Please note that most behaviour scales have not updated the description of behaviours to fit this example.

Behavioural expressions can often be a reaction to a changed environment or relationships with people; hence the importance of the relationship with care staff and why person centred care is so important.

### What are behavioural expressions?

The International Psychogeriatric Association (IPA) ‘Nurses Guide to Behavioral and Psychological Symptoms of Dementia’ provides a list of behavioural symptoms and psychological symptoms.

Behavioural symptoms	Agitation; Calling out; Crying; Physical aggression; Restlessness; Screaming; Sexual inappropriate behaviour; Verbal aggression; Wandering  These are usually identified by observation.
Psychological symptoms	Anxiety; Apathy; Delusions; Depressed mood; Disinhibition; Euphoria; Hallucinations; Misidentifications; Sleeplessness  These are usually identified by interview (with the person and caregiver), and some by observation.

Specific behavioural and psychological symptoms are not universally agreed upon in the literature as they are often interrelated, and they are not grouped for diagnosis purposes. It is not the grouping of behaviours that is primarily important, but it is important to understand the broad types of behaviour when reading the literature.

Many of the behavioural symptoms of BPSD relate to the wandering, verbal and physical behaviours targeted in the ACFI questions 7, 8 and 9.

Most of the psychological type symptoms of BPSD relate to mood and psychotic symptoms (e.g. confused thinking, delusions).

It is not of primary importance which group a behaviour or symptom belongs to. Rather it will be important to clearly describe the behaviour or symptom by what is seen or heard, without interpretation or judgement. This is the first step in unpacking the behaviour or symptom. Understanding of the person and the context of the behaviour is then vital in understanding the behaviour or symptom, and that requires looking at the interplay of physical, psychological, interpersonal, social and environmental factors.

### **Why is BPSD important?**

- With the aging of the world's population, there is a significant increase in the absolute number of older people with dementias
- Dementia is associated with progressive cognitive decline, a high prevalence of behavioural and psychological symptoms of dementia (BPSD) such as agitation, depression and psychosis
- BPSD are an integral part of the disease process and present severe problems to persons, their families and caregivers
- BPSD are treatable and generally respond better to therapy than other symptoms or syndromes of dementia
- Treatment of BPSD offers the best chance to alleviate suffering, reduce family burden, and lower societal costs in persons with dementia

Untreated BPSD contribute to:

- Early institutionalisation (Colerick and George, 1986; Morriss et al., 1990; Steele et al., 1990; O'Donnell et al., 1992; de Vugt et al., 2005);
- Increased financial cost (Cohen-Mansfield, 1995; Herrmann et al., 2006);
- Decreased quality of life for both the caregiver and the patient (Deimling and Bass, 1986; Burgio, 1996);
- Significant stress to caregivers (Rabins et al., 1982) and nursing staff in residential facilities (e.g., Rodney, 2000; Draper et al., 2000);
- Excess disability (Brody 1982; Hinton et al., 2008). Once symptoms are ameliorated or removed, functional level improves (reducing distress and improving quality of life).

*(International Psychogeriatric Association 'Specialists Guide to Behavioural and Psychological Symptoms of Dementia' Module 1, p.1.2-1.8)*

## *Behavioural Symptoms*

The following section is based on the *International Psychogeriatric Association 'Specialists Guide to Behavioural and Psychological Symptoms of Dementia' Module 2, p.2.11-2.13.*

### **Wandering**

This includes aimless wandering, exit seeking or trying to leave a building. It may be associated with faulty navigational ability (e.g., they feel lost), boredom or anxiety (e.g., looking for something or someone, don't know where they should be).

For example:

- looking for the bathroom
- looking for food or drink
- looking for a bed to lie down on
- acting on an old routine such as leaving work for home
- trying to escape the noise, tension or pandemonium of the environment
- searching for familiar faces (friends and relatives who may or may not be living)
- in pain, looking for relief
- desiring fresh air; may be triggered by outdoor clothing or view

### **Agitation**

Agitation (like BPSD) is not a diagnostic term, but rather a term used by clinicians for a group of symptoms that may reflect an underlying disorder. Cohen-Mansfield and Billing (1986), described agitated behaviour in three ways:

- It may be abusive or aggressive toward self or other
- It may be appropriate behaviour performed with inappropriate frequency, such as constantly asking questions
- It may be inappropriate according to social standards for the specific situation, as in taking off clothes in the activity room

Agitation usually increases with increasing cognitive impairment, and has been associated to a number of factors - neurobiological changes, health issues, pre-morbid personality, and the psychological and social environment. It is seen in physical symptoms (restless, wandering, hiding things etc) and verbal symptoms (constant requests, repetitive questions or complaints etc).

### **Aggression**

Verbal aggression such as screaming and cursing is often associated with depression, pain and other health problems. Physical aggression symptoms include hitting, pushing, biting etc.

### **Resistiveness to care**

Resistiveness to care is different to agitation; it is related to accepting assistance with daily activities like taking medicines, and undertaking ADLs. It is linked to the person's ability to understand the situation and what is expected, and is therefore likely to increase with the progression of the cognitive impairment, and it is often associated with behavioural expressions.

### **Inappropriate sexual behaviour**

Inappropriate sexual behaviour or sexual disinhibition may result in physical and/or verbal behaviours.

### **Sundowning**

This is BPSD in the afternoon and early evening. Dementia's can cause disturbance in the circadian rhythms which results in sleep disturbances. These symptoms have been also associated with hormonal changes (e.g. changes in body temperature), physiological and environmental factors.

### **Catastrophic reactions**

These are usually acute and brief and are usually an expression of overwhelming anxiety/frustration triggered by a task (e.g. getting dressed) or health factors (e.g., delirium, pain, infection, and certain medications).

## ***Psychological Symptoms***

The Cognitive Workbook discussed how depression and dementia are difficult to disentangle. Both share symptoms, for example, of being withdrawn, and depression is not uncommon in a person with dementia particularly in the early stages of dementia.

Depression symptoms can also include symptoms we typically associate with cognitive impairment (trouble thinking, concentrating, making decisions and remembering things).

Depression is a mood disorder that is much more severe than simply feeling down or flat emotionally after a loss or setback. It affects how a person feels, thinks, and acts and can lead to other emotional and physical problems. Symptoms are persistent day after day, and may include:



## Prepare

- Feelings of sadness, emptiness or unhappiness
- Outbursts of anger, irritability or frustration, even over small matters
- Loss of interest or pleasure in normal activities
- Sleep disturbances, including insomnia or sleeping too much
- Tiredness and lack of energy for even small tasks
- Changes in appetite —usually reduced appetite and weight loss, but cravings leading to weight gain in some people
- Anxiety, agitation or restlessness
- Slowed thinking, speaking or body movements
- Feelings of worthlessness or guilt, fixating on past failures or blaming yourself for things that are not your responsibility
- Trouble thinking, concentrating, making decisions and remembering things
- Frequent thoughts of death, suicidal thoughts, suicide attempts or suicide
- Unexplained physical problems, such as back pain or headaches

Many treatments are available, including medications and psychological therapies. There is evidence that exercise, getting outdoors and re-engaging in meaningful activities are all mood lifters. Depression is not a normal part of ageing, and people may not let others know they may have depression (BCOPE, p172).

In the most severe form of depression (psychotic) a person can have also have delusions (false beliefs, typically of being a bad person, deserving punishment or that bad things will happen) and hallucinations that may involve hearing voices, smelling unpleasant odours or other physical sensations (BCOPE, p172).

Early screening can assist in identification and timely intervention, and assessment by a mental health specialist is recommended for suspected depression and/or psychotic disorders (BCOPE, p172).

### *Psychiatric Symptoms*

Symptoms of psychosis include hallucinations, delusions, disorganised speech, abnormal psychotic behaviour, negative symptoms, impaired cognition, depression and mania. There are shared symptoms across psychotic disorders, cognitive impairment and depression. Many people with psychotic disorders also have cognitive impairments or depressive symptoms that affect their capacity to function (Diagnostic and Statistical Manual of Mental Disorders, 5<sup>th</sup> Edition, 2013). Following is a brief description of these symptoms.

Term	DSM-5 Description
Hallucinations	Perception like experiences without external stimulus. Occur in any sense modality, not occurring during falling asleep or waking up. (DSM-5)
Delusions	Fixed beliefs not amenable to change in light of conflicting evidence. Themes include persecutory, referential, somatic, religious, and grandiose. (DSM-5)
Disorganised speech	Inferred from the speech, answers to questions are obliquely related or unrelated, word salad is a severe example.
Abnormal psychotic behaviour	Rigid, bizarre posture, mutism, stupor (no motor response), excessive motor activity without cause, stereotyped movements, staring, grimacing, echoing of speech.
Negative symptoms	More common in schizophrenia, two most common are diminished emotional expression and avolition (lack of drive or motivation).
Impaired cognition	Complex attention (sustained, divided, selective, processing speed); Executive functioning (planning, decision making, mental flexibility); Learning and memory (immediate/recent/long term memory); Language (expressive and receptive); Perceptual motor (visual perception, visual constructional, perceptual motor, praxis, gnosis); Social cognition (recognition of emotions)
Depression	Depressed mood most of day, markedly diminished interest, significant weight change, insomnia or hypersomnia, psychomotor agitation/retardation, fatigue, feelings of worthlessness or hopelessness, diminished ability to think/ concentrate, recurrent thoughts of death, poor appetite/over eating.
Mania	Persistently elevated, expansive or irritable mood with goal-directed activity/energy.

### Hallucinations

The most common type of hallucinations in dementia are:

- Visual around 30%, up to 80% in person's with Lewy body dementia.
- Auditory around 10%
- Olfactory and tactile are rarer in dementia

Hallucinations may or may not be distressing to the person experiencing them, and may possibly be a visual misperception. Estimates of the frequency of hallucinations in people with dementia range from 12% to 49% (Swearer, 1994).

*(International Psychogeriatric Association 'Specialists Guide to Behavioral and Psychological Symptoms of Dementia' Module 2, p.2.6)*

## Misperception

Disorders of perception occur in the presence of external stimuli. There are 4 main types of misperception:

- Presence of persons in the person's own house (the 'phantom boarder' syndrome)
- Misidentification of the person's own self (often seen as not recognising their own mirror reflection)
- Misidentification of other persons
- Misidentification of events on television (the person imagines these events are occurring in real three-dimensional space)

An earlier study of misidentifications found them to be present in 16% of people with Alzheimer's disease and 17% of those with Dementia Lewy Body (Harciaiek & Kertesz, 2008). (*International Psychogeriatric Association 'Specialists Guide to Behavioral and Psychological Symptoms of Dementia' Module 2, p.2.7*)

## Delusions

The most common type of delusions in dementia are:

- People are stealing things
- Abandonment
- Home is not own home
- Spouse is an imposter
- Spouse infidelity

The frequency of delusions in people with dementia is cited as being between 10% and 73% depending on the study population and the definition of dementia (Wragg & Jeste, 1989). (*International Psychogeriatric Association 'Specialists Guide to Behavioral and Psychological Symptoms of Dementia' Module 2, p.2.5*).

## How Behavioural Expressions Interact with Other Domains

Behavioural and psychological symptoms may result from multiple causes although the actual origin remains undetermined. There are "neurobiological, psychological (premorbid personality features and responses to stress), and social (environmental change and caregiver factors) aspects (Kozman et al 2006 in [Hersch](#) et al 2007).

Therefore it is important to consider the impact of any other reduced functional abilities such as vision, hearing, mobility, incontinence etc. They may have other co-morbid health issues such as pain or depressive symptoms. [Hersch et al \(2007\)](#) indicates that


“Undiagnosed medical problems such as pain, depression, dehydration, sleeping difficulty, anxiety and delirium can all lead to agitation. (p.612)”

Resistiveness to care behaviours sometimes impact on the performance of daily living activities (ADLs). These activities vary from taking medicines to trying to assist the person with eating, personal hygiene and self care activities. In addition, as these are primarily private, intimate tasks, the person with dementia may feel unsettled and insecure when a carer attempts to undress them or help them with toileting. A verbal or physical behavioural expression may be a reaction to the fear.

A person with depressive symptoms also finds it difficult to engage in activities of daily living particularly morning hygiene activities, as their symptoms are typically worse in the morning. They may be withdrawn, feel too tired to participate, and lack interest in usual activities and social interactions.



## Identification Process



The steps in the process of **identifying** are:

- Gathering history from current documentation and information from carers, family and the consumer if possible;
- Identifying a need (e.g. initial nurse assessment/screen); and
- Completing a comprehensive assessment of the needs

## Gathering the History

What documents (before you start assessing) do you have which provide information on the resident you are focusing on? You will be able to build a picture of the person’s relevant history through current documentation (i.e. ACCR, medical notes) and their personal history through discussions with their family members and carers.

### Documentation

The current documentation should be reviewed, checking for relevant diagnoses, documented signs and symptoms, and previous assessments. Below is an example of the types of documents to be collated and reviewed and the information that is being sought for behavioural expressions.

Document	Looking for
ACCR	Part 4 (Q28): Mental and Behavioural Diagnoses; Part 5 (Q35) Assessment Summary ( cognitive behaviour/psychological symptoms) Part 5 (Q38 Q42) notes may inform on care strategies Parts 4/5: Notes on other impairments such as sensory loss that could further impact on behavioural/mood/psychiatric symptoms
Medical Notes/CMA	Mental and Behavioural Diagnoses, cognitive status, other impairments, behavioural concerns, mood, psychological and psychiatric symptoms, and assistance required
Clinical Report on Depression	Mental and Behavioural Diagnoses, mood symptoms, cognitive behaviour/psychological aspects, comments and care strategies

## Behavioural Expressions Topic



Mental Health Professional notes/assessment	Mental and Behavioural Diagnoses, notes on symptoms and strategies

Here are a few examples of how the collated diagnoses and issues may be used to understand the person's care needs:

Information	Impact
<p>Diagnosis or evidence of</p> <ul style="list-style-type: none"> <li>• cognitive impairment</li> <li>• behavioural concerns</li> <li>• mood symptoms</li> <li>• psychosis symptoms</li> </ul>	<p>An accurate diagnosis is essential for understanding the behavioural expressions and for developing appropriate strategies and interventions.</p>
<p>Specifics about the behaviour and symptoms</p>	<p>It is important that the behaviour or symptom is objectively described e.g. what is seen or heard by staff. This will assist staff communication, provide the most accurate information and ensure the Behaviour Records are accurate. It will also assist in identifying evidence-based care strategies.</p>
<p>Behavioural and Psychological Symptoms of Dementia</p>	<p>The person with dementia has increasing trouble deciphering and interacting with their environment</p> <ul style="list-style-type: none"> <li>- ability to communicate their needs</li> <li>- ability to understand verbal requests</li> <li>- ability to remember how to do basic activities (e.g. affects motor skills)</li> <li>- misinterpreting their environment (e.g. hearing and seeing things, holding incorrect beliefs)</li> </ul> <p>They can become physically and verbally restless and aggressive due to the confusion they experience, they may not understand where they are, why they are there or what they should be doing etc.</p>
<p>Mood symptoms</p>	<p>Depression has been under diagnosed in residential care. It is important to increase the awareness of mood symptoms as mood affects participation in everyday activities. Residents may be withdrawn, sad, or display symptoms of cognitive impairment (e.g. memory loss). There are both non pharmacological and pharmacological treatment options with proven efficacy.</p>
<p>Sensory Loss(e.g. sight, hearing)</p>	<p>Most likely to increase the affect of the cognitive/behavioural impairment, making it more difficult for the person to participate in activities, may become withdrawn.</p>



## *Resident and Family*

It is important to seek input from the resident and his or her family members. The family knowledge of the person's medical and personal history, cognitive functioning, communication style and preferences, and routines is an important source of ongoing information for residential care staff. Families and carers who are regular visitors may be more likely to notice the subtle changes in cognition and behaviour which can serve to highlight the need for further investigation. Family members may also know helpful hints when supporting a person with dementia. For example the conversation cues that can help to re-settle them (i.e. significant life events, pleasant events, favourite pastimes, work history, personal history).

## Screening

For a new resident, staff should note and document (in progress notes and against a standardised checklist e.g., the modified Behavioural Assessment Form and the Behavioural Descriptions list) all disruptive behavioural symptoms expressed by the person. Mood may not appear disruptive but it is likely to be concerning to the resident themselves. Behaviours and symptoms may be disruptive or concerning to the resident as well as other residents and visitors or staff.

When in the screening phase staff should be observing the person to identify if behaviours are present. **The behaviour may or may not be associated to an ACFI behaviour item or question at this stage. The aim at this point is NOT about completing ACFI documentation.** Staff should document these interactions providing an objective and individualised description of the behaviour, including what was occurring at the time, the type of staff response and the outcomes. At this point the resident may still be settling in, (a change of environment is an unsettling experience for anyone and especially for a person with a cognitive impairment), so it is best not to do the ACFI assessments in the first week.

The most important thing at this stage is to be developing a description of the behaviour/symptom, based on what is *“seen or heard”* by staff or their family.

- Staff should document what is seen and heard including the contextual information.
- Staff teams should then identify which behaviours are problematic (i.e. require staff intervention). Sources of information that may be considered could include:
  - Gathered documents (admission documents, medical notes etc)
  - Interviews with family
  - Nursing notes, handover notes etc
  - Behavioural Description List



### The Behavioural Descriptions List

Discuss “*what is seen and heard*” with other staff and ensure there is an agreed description for consistent staff communication and assessment. After the behaviour has been uniquely described, the behaviour can then be identified in the Behavioural Descriptions List (a copy is provided in the Behavioural Expressions Assessment Pack). This list of behaviours and symptoms covers the ACFI behaviours, Cornell Scale for Depression in Dementia and psychiatric symptoms.

Provide each behaviour or symptom with a unique description based on what is seen and or heard. Only behavioural symptoms that require staff intervention should be considered. Not all behavioural symptoms will map to an ACFI question, however all behavioural expressions that cause distress or disruption should be investigated and managed. The behavioural expressions management approach is a comprehensive care planning approach driven by quality care beyond compliance and ACFI needs.

There are a limited number of screen items covering mood symptoms for older people in the CHA, if used for the initial nurse assessment. To screen across all symptoms, it is recommended that the Behavioural Descriptions List be completed (a copy is found in the Behavioural Expressions Assessment Pack). Table 2 provides a modified extract from the Behavioural Description List. This snapshot covers some of the behavioural symptoms and demonstrates how the behavioural expression can be mapped to the ACFI questions. In the full version the list covers all symptoms and maps to all ACFI questions:

- Behavioural symptoms- some which map to Wandering (ACFI 7), Verbal Behaviour (ACFI 8) and Physical Behaviour (ACFI 9)
- Psychological symptoms - which map to Depression (ACFI 10)
- Psychiatric symptoms

Note that an individualised description should be provided of the behaviour or symptom of interest as well as documenting in the progress notes during this investigation period. It is important that there is a clear understanding across the care team of the behavioural expressions, the full descriptions of the behaviour will provide that picture.

The screening stage should be undertaken over the first two weeks which allows for new residents to settle in before starting ACFI assessments. The care team will obtain enough information to identify the behaviours and symptoms that will be targeted in the next step of the comprehensive assessment. Figure 2 provides an overview of the assessment process the screening step is undertaken in week 1, and the comprehensive assessment steps in weeks 2-4.




**Table 2: Modified extract from the Behavioural Descriptions List**

BEHAVIOURAL DESCRIPTIONS LIST	MAPPING			
	ACFI 7	ACFI 8	ACFI 9	ACFI 10
<i>Must require staff intervention. Report what you see or hear.</i>				
Interfering with other people or their belongings while wandering	W1: Interfering while wandering		P2: Socially inappropriate handling of objects	
Trying to exit the building or property <b>e.g., Mr. Grey was pacing around the facility and garden when he was observed trying to climb over the fence</b>	W2: Trying to get to inappropriate places			
Trying to get into locked areas, offices, other resident's rooms	W2: Trying to get to inappropriate places			
Aimless wandering not disturbing others	n/a	n/a	n/a	n/a
Verbal refusal to participate in daily care- dressing, washing, hygiene, medications <b>e.g., Mr. Grey refused to take his underwear off during his morning shower, he would scream out “no, no, leave them on” and threatened to “knock their blocks off”, if the staff tried to pull them down.</b>		V1: Verbal refusal of care		
Cursing, swearing, temper outbursts, verbal anger, verbal combativeness, screaming		V2: Verbal disruption to others (not an unmet need)		Irritability: easily annoyed, short tempered, impatient, angry
Verbal accusations based on excessive/unrealistic suspiciousness		V3: Paranoid ideation that disturbs others		
Verbal sexual propositions, sexual innuendo, sexually threatening language		V4: Verbally sexually inappropriate		



Figure 2: Behavioural Assessment Process





## Comprehensive Assessment

Following a positive identification through the screening phase a behavioural symptom will have been identified as being present, and requiring staff intervention. The following approach is recommended and the assessment tools listed below can be found in the Behavioural Expressions Assessment Pack:

- Modified Behavioural Assessment Form (BAF) for behaviour symptoms
- Modified Cornell Scale for Depression (CSDD)

The next step is to determine if it is a 'behavioural' or 'mood' or 'psychiatric' symptom. The Mapping Document found in the Behavioural Expressions Assessment Pack will assist staff to identify what grouping the behaviour or symptom is associated with. , A brief overview is as follows:

### Behaviour

- Wandering behaviour (e.g. interfering while wandering, exiting behaviour, trying to get to inappropriate places, aimless wandering, pacing)
- Physical behaviour (e.g. harming self/others/property, disrobing, inappropriate sexual behaviour, eating or drinking inappropriate substances, hiding things, repetitive movements, constant restlessness, )
- Verbal behaviour (e.g. verbal refusal to participate in ADLs, strange noises, negativism, repetitive sentences, cursing, screaming, excessive suspiciousness or verbal accusations of delusional thoughts)

### Mood/depression

An expression of mood/depression could be:

- Mood symptoms (e.g. anxious expression, crying, lack of reactivity, irritability)
- Behavioural disturbance (e.g. agitation, displayed in pulling one's own hair or picking at clothing/ physical restlessness etc, slow movements, loss of interest)
- Physical signs (e.g. appetite, weight loss, lack of energy)
- Cyclic functions (worse in morning, sleep difficulties)
- Ideational disturbance (e.g. feels worthless/a failure, expects the worse)



### **Psychiatric**

Residents with a neuro-cognitive psychiatric disorder (e.g. delusions, hallucinations) will likely exhibit a behaviour that is described by the symptoms outlined above under the behaviour heading.

For example a resident with a delusional disorder may (i) start to act out on their irrational belief verbally or even physically which ultimately becomes disturbing to other residents and staff or (ii) they become extremely bothered by the irrational belief which causes a major disruption to their quality of life. It will not usually be the 'delusion' that staff become aware of initially, but rather a disturbing behavioural 'acting out' due to the underlying delusion e.g. verbal disruption or physical behaviour that can be subsequently linked back to the delusional disorder.

For these psychotic symptoms, you should still describe the symptom/s objectively (i.e., what you see and hear the person doing), and refer the person to a Mental Health specialist for further assessment. It is the severity of the symptoms in psychosis which will guide the likely course and intervention required (DSM V, 2013).



## *For wandering, verbal or physical behaviour*

The suggested approach is:

- Modified Behavioural Assessment Forms is undertaken for 7 days. The data can then be transcribed to the ACFI Behaviour Records.
- The modified Behavioural Assessment Form (found in the ACFI Assessment Pack) will record the detailed behaviour description, frequency and pattern of the behaviour, and the contextual information. It is important that the link from the objective description as identified during screening is continued into the comprehensive assessment step. A further description of this tool is provided below.
- The description of the behaviour will help to determine (i) is it of concern to the person and/or to others (ii) does it need a referral and (iii) what strategy would be appropriate to assist settle or manage the behaviour.
- Complete the modified BAF and the Behaviour Record by transcribing the behaviour type and frequency data, this comprehensive information will inform on possible strategies and complete the ACFI ratings.

### **The Modified Behavioural Assessment Form**

The modified Behavioural Assessment Form (BAF) provides a standardised way to describe the contextual setting (a copy is found in the Behavioural Expressions Assessment Pack). The Behavioural Assessment Form (BAF) has been modified to include standardised responses. The BAF is based on the ABC theory of behaviour, an approach supported by the literature.

*“Behaviour does not occur in a vacuum. Whether we can see it or not, there is always something that starts it off. Sometimes this is an event such as the delivery of a meal; sometimes it is a thought such as ‘I am missing my husband.’ The trigger to the behaviour is called the **Antecedent**; it is the A of the ABC.*

*The **Behaviour** itself is the B and the **Consequences** of the behaviour are the C. The consequences have a strong influence on whether or not the behaviour will be repeated. If the consequences are experienced as rewarding the behaviour is likely to go on or be repeated, if they are not rewarding the behaviour it is likely to stop, eventually.*

*Effective behaviour management begins by trying to clearly describe the behaviour, identifying the triggers and analysing the consequences to see what is keeping it going. In other words applying the ABC approach.”(Behaviour Assessment Form, NATFRAME pp. 145)*



Behaviour assessment and management is not well understood or documented in residential aged care. To address this gap the EBCAT recommendations provide a process that produces:

- Accurate behaviour descriptions based on what is seen and heard
- Documentation of the contextual information (i.e. the circumstances at the time of the event) to ensure the behaviour is fully described
- A relevant documentation base that will then (i) allow the development of evidence informed strategies and interventions and (ii) also provide helpful information if external referrals are indicated

It is important that staff are encouraged to describe the behaviour in its natural context not by the ACFI broad behaviour descriptions during both the screening and the comprehensive assessment steps.

The modified BAF (a copy is found in the Behavioural Expressions Assessment Pack) has been recommended, because it collects both frequency and standardised contextual information. The modified BAF is not commenced until week two as the frequency data for the ACFI Behaviour Record cannot be started till after the first week to allow for new residents to settle in. The use of the modified BAF in the comprehensive assessment step will provide data that can be transcribed to the ACFI Behaviour Record and provides the detailed contextual information required for developing interventions and care planning.

The modified Behaviour Assessment Form items cover:

Questions	Responses
<b>Antecedents</b>	
Date and time, How long did it last?	
Place	<ol style="list-style-type: none"> <li>1. Travelling off campus</li> <li>2. Outside of building (but within the grounds)</li> <li>3. Bedroom</li> <li>4. Ensuite</li> <li>5. Shared bathroom</li> <li>6. Corridor</li> <li>7. Dining area</li> <li>8. Activity area</li> <li>9. Other (describe)</li> </ol>
Who was there?	<ol style="list-style-type: none"> <li>1. Other resident/s</li> <li>2. Own visitor/s</li> <li>3. Other visitor/s</li> <li>4. Doctor</li> <li>5. Care staff (describe)</li> </ol>

## Behavioural Expressions Topic



Questions	Responses
	<ol style="list-style-type: none"> <li>6. Therapy staff</li> <li>7. Activity staff</li> <li>8. Domestic staff</li> <li>9. Maintenance/gardening</li> <li>10. Other (describe)</li> </ol>
What interaction was going on?	<ol style="list-style-type: none"> <li>1. Self care (nutrition/walking/personal hygiene/toileting, self medicating)</li> <li>2. assisted care (nutrition/walking/personal hygiene/ toileting, medication round, complex health care)</li> <li>3. Medical treatment/visit</li> <li>4. Therapy (physio, massage, describe other)</li> <li>5. Activities</li> <li>6. Other (describe)</li> </ol>
What else was happening?	Describe the environment e.g. Noise, stimuli, entertainment, TV, lot of people etc.
<b>Behaviours</b>	
Observed Behaviour	Describe what was <b>seen and heard</b> <i>For example: Mr. Grey was pacing around the facility and garden when he was observed trying to climb over the fence</i>
<b>Consequences</b>	
Describe the intervention	How many staff, what is the intervention, other resources required, time taken
Disruption Level	Rate the disruption <ul style="list-style-type: none"> <li><input type="checkbox"/> (0) None (Not disruptive no intervention by staff)</li> <li><input type="checkbox"/> (1) Mild (A little disruptive co-operative response to intervention, not disruptive to other residents or visitors)</li> <li><input type="checkbox"/> (2) Moderate (Moderately disruptive not always co-operative, but can be resolved with intervention, sometimes disruptive to other residents or visitors)</li> <li><input type="checkbox"/> (3) Severe (Very disruptive sometimes requires immediate intervention, interferes with others, their belongings or visitors, asocial behaviour)</li> <li><input type="checkbox"/> (4) Extreme (Extremely disruptive always requires immediate intervention, wakes others at night, disruptive to others during the day, requires one or more staff, attention or constant attention)</li> </ul>
Rate the effectiveness of the intervention	Effectiveness is rated <ul style="list-style-type: none"> <li><input type="checkbox"/> (0) No intervention by staff</li> <li><input type="checkbox"/> (1) Very successful, co-operative response, settled quickly</li> <li><input type="checkbox"/> (2) Successful, settled within 5 minutes</li> <li><input type="checkbox"/> (3) Somewhat successful, settled for short time only or requires constant attention</li> </ul>



### *For a mood symptom*

The mandated approach is:

- The modified Cornell Scale for Depression (CSDD). A copy is found in the ACFI Assessment Pack and the ACFI User Guide provides detailed instructions on how to complete this assessment.

The ACFI Assessment Pack provides the modified CSDD tool which is a validated (objective and standardised) assessment for identifying those residents with symptoms of depression. The CSDD assessment is:

- Based on signs and symptoms that staff have opportunity to notice, it does not include symptoms such as phobias that require a reliable self report
- Has defined signs and symptoms so as to reduce the subjective interpretation
- Appropriate and validated for aged care staff that have been appropriately trained on the application of the tool
- A screening test and requires GP/specialist follow-up for diagnosis and management assistance (following good practice)
- A tool that uses multiple sources of information to complete which improves the objectivity, accuracy and quality of information collected
- A tool that fosters staff awareness of signs and symptoms of depression
- A tool that can be used to objectively assess for improvement

The modified CSDD assessment in the ACFI Assessment Pack provides detailed process steps to ensure an accurate assessment is completed. For example the modified CSDD instructions indicate:

- The symptoms must have occurred in the past week (and should have a description of the symptom/behaviour in context for that resident, not a generic description)
- The exclusions due to dementia etc
- The descriptions of how to rate the severity of symptoms such as mild or severe effects (based on frequency and minor/major interference):-





- Mild = minor interference (regular encouragement required with ADLs, social and interpersonal) and symptoms occur occasionally (not every day)/or often (nearly every day)
- Severe = major interference (limited participation in ADLs, social and interpersonal) and symptoms occur often

The modified CSDD criteria guidelines are recorded in the sub-steps. It is important to indicate the response to the sub steps in each question of the assessment and not just provide the score for each item. **This is a valuable source of information that can inform on strategies that are important regardless of the final summary score.**

If a resident is identified as having a possible change in their behaviour or mood status, it is recommended that this assessment phase is repeated.

## Clinical Risks

If depression is identified, it is recommended that the Standardised Care Process (SCP) Depression be commenced (a copy is found in the Behavioural Expressions Assessment Pack).

The Depression SCP describes a systematic approach to addressing depression:

- a. A recognition and assessment step starting at admission or as required, using the Cornell Scale for Depression
- b. Interview resident/family for history
- c. Monitor resident
- d. Repeat assessments for depression and associated impairments as required or every three months (CSD,PAS-CSI, Functional ability)
- e. Referrals to GP for diagnosis, psychiatric services for severe symptoms
- f. Develop individualised care plan, considering lifestyle, exercise, pastoral care
- g. Monitor the effectiveness and symptoms
- h. Provide information about depression to resident/family
- i. Staff knowledge and education requirements – causes, symptoms and levels of depression, treatment options

Physical Restraint is not considered good practice, if there is a clinical risk of physical restraint, the SCP Physical Restraint is recommended as it offers alternatives (a copy is found in the Behavioural Expressions Assessment Pack).

Prepare

Identify

### Bringing the information together

The gathering and review of documents followed by the assessment of behaviours and mood should include information from other domains in order to avoid missing relevant information. These might include:

- Sensory impairment (e.g. as identified in an initial nurse assessment)
- Physical impairment (e.g. as identified in an initial nurse assessment)
- Medications being taken as reported in the medication record or chart

### Back to Mr. Grey

*Mr. Grey has been at the home for a week now. He arrived at the home in the company of his son and wife. John had convinced Mr. Grey that the move was necessary because of Mrs. Grey's health problems, but Mr. Grey seemed quite down and withdrawn.*

*In spite of his mixture of vascular dementia and Alzheimer's disease, Mr. Grey was relatively healthy.*

*The staff reported that while the personal history they learnt from Mrs. Grey was useful in gaining his cooperation during ADLs (talking about musicals and his plumbing job), he is not cooperative with changing his underwear. He constantly paced around the facility and did not interact with the staff or residents. The home was securely fenced but six times in the week Mr. Grey was found trying to climb the fence.*

*To address the wandering behaviour:*

*(i) The dietitian recommended a supplement to fortify his diet due to his 'restlessness' and (ii) the physiotherapist introduced an exercise program to provide a controlled outlet for that restlessness and (iii) the psychogeriatrician recommended an anti-depressant, in part to reduce Mr. Grey's anxiety and restless behaviour, and in part to lift his apathetic mood.*

*The staff realised that they needed to know what might set off his wandering behaviour (the antecedent). What was the trigger that was making Mr. Grey think he had to leave?*


*They started a Behaviour Record: There was a set pattern- he set off in the morning (about 8-9am) or in the afternoon (about 3pm).*

*They watched more closely at these times (completing a modified Behaviour Assessment Form) and found a delivery van's often turned up at the home in the morning, and this was visible to Mr. Grey if he was in the garden. Was he reacting to old routines – being picked up for work after breakfast?*

*In the afternoon at 3pm the next care shift started, there was a lot more activity with people coming and going. Did he think this was a cue to go home?*



## Implementation Phase



This phase of the toolkit demonstrates how to use the information gained from the identification phase to implement interventions and care planning.

The implementation process has three main aspects. These are:

- Undertaking **referrals as identified** to gather further needed information;
- Designing **interventions** and developing **care plans that provide a coherent picture of what is to be done and why**;
- Completing **other documents** that support the care process and the accountability systems, such as for the ACFI funding claims and quality of care aspects for accreditation

## Referrals

If there is an identified need for further behaviour assessment or management that cannot be provided on site, a referral to a Dementia Behaviour Management Assessment Service (i.e., DBMAS or CDAMS) or a dementia consultant could be considered. If there is an identified need for further mood assessment or management that cannot be provided on site, then a referral to an Aged Psychiatry service or private psychiatrist or psychologist could be considered.

If there is an identified need for medication or medical specialist, then the matter should be directed to a Medical Practitioner (i.e. GP), who can implement treatments, or make referrals to medical specialists such as the geriatrician or psychogeriatrician.

We recommend that your home develops referral lists, based on what your health service can provide, what is available in your local area and a list of important website contacts to find a practitioner or advice, for example (Table 3).

After completing the behavioural screens and assessments (modified BAF, Behavioural Description List, and the CSDD if appropriate) there will be informative, objective and detailed information suitable to complete a standardised referral template to behaviour support services.

**Table 3: Referral for Behavioural Expressions**

Health Professional	Source	Contact
Medical Practitioner	Health Service X	Name, contact details
Geriatrician	Health Service X	Name, contact details
Psychogeriatrician	Health Service X	Name, contact details
Aged Psychiatry Services	Health Service X	Name, contact details
Behavioural Assessment Team	Health Service X	Name, contact details
Dementia Consultant	Health Service X	Name, contact details
DBMAS	Health Service X	Name, contact details

## Interventions

In the identification phase, the residents' healthcare and personal needs were identified. The intervention program will address these identified issues, by developing strategies to improve or maintain the residents' health status and their quality of life.

The intervention program targeting behavioural expressions should involve evidence informed strategies (refer following) that address the behavioural needs, and importantly the strategies should be specifically tailored for each individual to be most effective. The approach and actions you use to support one resident may be very different from those implemented for another resident.

When designing interventions consider the resident history and personal preferences, the assessment outcomes, the context the strategy will operate in (i.e. the physical environment, the social environment), the knowledge and attitudes of staff, residents and family, and the types of resources required and their availability. Interventions are likely to be medical, psychosocial, educational or nursing in nature.

It is also important that all staff follow a systematic process when implementing an intervention 'program'. This will increase the likelihood of your intervention's success. Having a systematic process which you can describe will also enable other staff to repeat your interventions if they prove successful.

Interventions may be based on pharmacological or non-pharmacological approaches. A Pharmacological approaches should be developed by a Medical Practitioner, the resident's GP or via a referral to a medical specialist e.g., geriatrician, psychogeriatrician or a dementia specialist service.



## *Non Pharmacological Approaches*

Good dementia care can be good care for any resident. Interventions that include a relaxed social environment, access to meaningful activities, positive staff- resident relationships should be considered for all residents. Here are some examples of non pharmacological approaches.

Physical Environment	The physical environment should be calm and supportive- small, homelike, with ready access to staff, and an arrangement of shared areas, single bedrooms with ensuites and access to safe garden walks. Noise and lighting should also be considered.
Relationships	Staff, other residents and the community should engage with the person as far as possible and encourage social interaction.
Person Centred Care	Taking into account the cultural, family values, interests and likes of the person, and understanding the person. Looking after the whole of the person, and make sure their health needs are attended to.
Activity and recreation	Meaningful activities that the person can relate to, even if they have memory loss.
1-1 interaction	Engaging the person in their surroundings, in conversation. Using daily interactions to engage the person. Ensuring the work place practices (roster, attitudes) support staff to interact with residents.
Physical activity	Exercise, incidental activity is good for the body and the brain.
Pleasant Events	Provides an opportunity to improve or maintain mood.
Sleep	Allowing people to sleep in and awake naturally.

When a person with dementia has a limited vocabulary, the behaviours may be expressing “I want to go home” or “I’m uncomfortable, I need some reassurance”. The key is to decipher the spoken and unspoken message by having a background history on the person - the history we have collected in the identification phase, especially the personal family, previous interests and working life.

Sometimes reminiscence can be comforting. “Do you mean you want to go back to Bendigo, where you grew up? Tell me what it was like there.”



When a man is worried about finding his spouse, for example, sometimes simply giving him a chance to reminisce about her is comforting. If his wife has died, a picture of her can help him reminisce. Perhaps a tape or video recording made by his wife or a family member will offer a reassurance and can be played anytime.

## *Communicating Effectively*

When we are talking about behavioural expressions as a form of communication resulting from a cognitive impairment and a diminished verbal vocabulary, it is doubly important that care staff verbal and non-verbal messages to people are clear and sensitive.

Some suggestions for providing verbal messages to people with cognitive impairment are:

**Don't underestimate what people understand.** Always include them in conversations if possible.

**Introduce yourself when you come into the person's presence.**

**Slow down and be as patient as possible.** Slowing down is probably the most important thing we can do in communicating with anyone and especially someone with a cognitive impairment or impaired mood.

**When giving directions, use short sentences, and take one step at a time.** "I'm going to help you put on your blouse. Let's start with this sleeve."

**Don't over-explain.** As dementia progresses, the people who have it tend to speak less. We have a tendency to fill in those conversational gaps with detailed explanations which can make the person more anxious.

**State things positively.** If you say to a person, "Don't go outside," he is likely to focus on going outside. Avoid negatives and say what you really want, i.e. "Stay inside."

**Issue an invitation.** If you ask the person with dementia if they want to do exercises it will likely result in a "No, thank you". Instead, issue an invitation such as "Mr. Grey, would you please come with me to the sing-along? I need your wonderful voice"

**Use concrete terms.** "Put that over there" is likely to be meaningless to the person with dementia. "Please put your plate on the table" is better.

**Respect the person's right to say "no."** When things aren't working, leave, and come back later.



**As a person’s dementia progresses, avoid open-ended questions and limit choices.** Asking a person with dementia, “What would you like for lunch?” may be overwhelming. Asking, “Would you like the cheese or the tuna sandwich?” is an easier decision.

**Use multisensory clues.** A bath towel and soap can indicate that it is time to have a shower. The more ways you can signal what is happening next, the more likely it is that the person with dementia will understand you and start thinking about the next step.

**Having a conversation.** If you ask multiple questions in a row, the person with dementia may become more confused. Ask one thing at a time, for example:

- Where you go to school?
- Did you walk to school?
- Was it far away?
- Did you like your teachers?

**Substitute statements for questions.** For example if you ask while walking with Mr. Grey, “Mr. Grey, would you like to use the bathroom?” he will likely say no, but if you say “here is the bathroom” it sets an expectation and may get a different response.

**Be silent when necessary.** Some people with dementia need all their powers of concentration to focus on the task you are assisting them with. If you start to talk about the weather, they will forget they were trying to put on a shirt or blouse, and be confused for a while before they can return to the task. It is important to recognise what an individual resident needs in any given situation to make them as comfortable as possible.

Back to Mr. Grey

*For Mr. Grey, the harder challenge is to solve the “man on a mission” sort of wandering that he frequently does. This is sometimes referred to as “exit-seeking” behavior because the person is determined to leave the building or property. The feeling of having obligations – work to get to, children to pick up, dinner to prepare – is often strong in the middle stage of Alzheimer’s disease. If you try to stop mission driven people, they may become frustrated or anxious because you don’t allow them to complete their tasks.*

*Sometimes, as happened when Mr. Grey lived at home, letting him go, and counting on him to return on his own when the urgency left him, can work just fine, at least for a while. That isn’t usually practical and it may not be safe in many circumstances.*

*Sometimes a person can be distracted by a conversation (especially over a cup of tea and biscuits). For example staff could ask him about his work, “Was it hard, or messy? Were the customers pleasant?” Or ask for his opinion. Mr. Grey may also retain his career-related vocabulary. So ask, “My kitchen tap is old and has started to drip. Do you think I should replace it?”*



*Sometimes a person can be distracted by a job to do. It may be slightly related or totally unrelated to his occupation or skills. Sometimes you need to be creative. Placing an item of interest in a garden may provide the distraction that reduces the urge to leave the home. It is important to include staff and family members when developing strategies and to see what works best for that resident and the facility.*

*The staff have identified and described the wandering behavior, completed a further behavior assessment looking at the antecedents and consequences of these occurrences over the last week, and talked with the family.*

Following is a sample item from Mr. Grey's now revised care plan related to his wandering behaviour.

*Staff planned new routines and distractions for the two periods that set off the exit seeking behaviour.*

*There were two secure gardens at the home and the second garden area was directly off the dining area and did not give a view of the delivery vehicles. The activities staff set up an after breakfast activity just off the dining area where Mr. Grey had breakfast. He (and other residents) were invited to listen and sing along to resident's favourite music for one hour. This activity was attractive to Mr. Grey's personal interests and distracted him away from the other garden area in the morning. It was also hoped that this pleasant event would encourage interaction between Mr. Grey and some other residents. Another goal was to establish a new morning routine for Mr. Grey.*

*A new routine was also needed in the afternoon about the time that many staff left. They asked Mrs. Grey to visit in the afternoon when possible. From talking with Mrs. Grey they learnt that Mr. Grey had enjoyed shredding their papers and taking them to the local council dog pound. It was decided that on two afternoons, they would start shredding paper and cardboard for the local dog pound.*

*It was hoped that by distracting Mr. Grey with useful work during the two times he was most likely to try and leave, he would stop doing so.*

*It was important that Mr. Grey's and other resident's access to the external areas was not affected, as it was good for all resident's to get outdoors for walks in the secure garden area. They would also see if they might be able to get him outdoors for some gardening activities – another way to give him fresh air and a sense of purpose. They would re-evaluate how things were going a week later.*





## Care Planning

A comprehensive care plan will be more than a summary of care needs, it will be individualised for each resident and reflect an understanding of their story, history, preferences and care needs.

Facilities may have their own care plan or the NATFRAME Care Profile could be used as a starting point e.g. as a summary care plan. Although the NATFRAME Care Profile will probably not record enough details for your home, it is an example of how to collect and document information in a systematic, professional and accountable manner.

## Goal Setting

Moving beyond compliance, the care planning should also address setting goals based on clinical outcomes and the resident's perspective on their care needs and what is important to maintain their quality of life. It is recommended that a **Quality Of Life (QoL)** questionnaire be used to ascertain the resident's view of their quality of life. This could be implemented during the resident/family meeting to sign off on the Care Plan. The goals developed with the resident/family should cover **resident QoL, clinical outcomes and quality care outcomes** to address a wide range of documentation needs (i.e. consumer satisfaction and clinical expectations).

**SMART Goals** are an example of a standardised approach to goal setting with measurable outcomes. The SMART acronym ([www.projectsmart.co.uk](http://www.projectsmart.co.uk)) stands for goals that are:

**Specific**, that is, they provide clarity, focus and direction.

**Measurable** - Objective measures can demonstrate the effectiveness of the goals.

**Action-oriented**, that is, they provide a strategy for achieving them.

**Realistic** – because if they're not, we're just setting up for almost certain failure that will then impact on the residents motivation, interest and involvement; and

**Time-based**- meaning for a specific duration. That enhances their ability to be measured and re-evaluated on an ongoing basis

## Role of Documentation

Documentation of care is essential because members of the care staff are accountable to each resident for their actions and decisions. Accountability is also a vital function of professional, legal, accreditation and funding mechanisms and processes.



Documentation is also a communication tool about what has been investigated and the information collected, what interventions have been tried for a particular problem with a particular resident and to identify what is and isn't working. It's a communication tool that can help all staff give the best possible care.

Good documentation supports the nursing process which drives the care. It should incorporate evidence informed care, evidence-based assessments and interventions, utilise staff skills and drive a quality care approach.

The documentation process should leave a trail that provides robust and accountable information, leaving your facility audit ready. The documentation is the basis of an evidence profile which provides objective and transparent information for the reason that assistance is or is not required. This supports the ACFI claim which is an outcome of the process, not a driver.

The process for documenting care is as follows:

- Summarise the relevant points from the collated background information and personal history
- Record the assessment and referral outcomes
- Record the identified capabilities and need for assistance following an analysis of the background information and assessment outcomes. This forms the basis of an evidence profile which provides objective and transparent information about the reason that assistance is or is not required.
- Prepare the care plan with details on the care to be provided, why, and the resident's goals and desired outcomes (in consultation with the family if appropriate). Record the evaluation of the care provided and the resident's response to the care and quality of life outcomes
- Continue to document new activities or identified issues (e.g. developed strategies and changes to care planning) in the progress notes and in the care plan, as this assists the information and communication flow and allows the process to be tracked for quality purposes.
- Incident forms should be completed recording episodes of falls. The analysis of such incidents should take into consideration the resident's right to take risks. For example, a resident who is capable of making his own decisions may choose to walk outside even though he cannot be fully supervised at all times. This would be documented in the resident's goals and care plan



## Linking the Evidence

This is about showing the link between the evidence from the background and assessment information (what has been collected in the identification phase) and the care that has been provided to address the identified care needs (in the implementation phase).

The links are:

- Begin by looking at and noting any underlying issues (e.g. dementia diagnosis) or symptoms (e.g. increasing agitation in the afternoon), connect the link to the body structures and/or functions that are impacted (e.g. disruptive behavioural symptom)
- Describe the associated activity limitations (e.g. requires supervision in afternoon). It's important to look at remaining strengths (e.g. enjoys being helpful and busy)
- Describe the intervention program and how it addresses the limitations to assist the resident to improve or maintain his or her status and ensure continued participation in a quality life (e.g. introduce new activities in the afternoon)
- Finally, define the care goals (to reduce afternoon agitation) and resident goals (feeling useful) that will be evaluated. Goals give purpose to the care and provide a measurement for evaluating the outcomes.

Below is an example of linking the evidence from the case study of Mr Grey:

- There was a diagnosis of moderate to severe dementia, with symptoms of excessive wandering and exit seeking behaviour. Also symptoms of verbal combativeness to his wife. Depressed mood was noted after admission to the home. This was evidenced in the Medical Notes, ACCR, and progress notes.
- From the identification process, we identified the body structures/ functions that were affected. This was mood, behaviours and cognition. The cognitive assessment showed concentration issues, short term memory loss and disorientation to place and time. The Behavioural Expression assessments showed physical restlessness, exiting behaviour, verbal refusal of some personal aspects of care, and loss of interest. These issues have resulted in reduced independence in personal hygiene and limited interaction with staff and residents
- Interventions included distraction combining pleasant events, personal hobbies and interests i.e., music activity after breakfast, and paper recycling twice a week.
- The aims of the care plan were then defined as goals (e.g. reduce exit seeking behaviour, increase social interaction) that will be evaluated.



## Completing the ACFI documentation

The data collected can now be used to complete the **ACFI 7, ACFI 8, ACFI 9 and ACFI 10 checklists and ratings** (Tables 4 and 5).

The mandatory ACFI assessments for ACFI 7, 8, 9 and 10 (i.e. Behaviour Records and the Cornell Scale for Depression in Dementia) should be no more than 6 months old and should reflect the current needs of the resident.

To support a B, C or D rating in ACFI 7, 8 and 9, the required ACFI assessment is the Behaviour Record (Table 4). It is recommended that users refer to the **ACFI User Guide and Assessment Pack** to support implementation of the tool. The ACFI claim is directly linked to the frequency as recorded in the Behaviour Record. It is the same approach for all three behaviour groups. The mapping shown on the Behavioural Descriptions List (found in the Behavioural Expressions Assessment Pack) will assist you to determine which ACFI claim(s) to make.

**Table 4: ACFI Questions 7 to 9 Checklists**

ACFI 7, 8 & 9 Behaviour Checklists	Rating	Where to find the supporting information
Behaviour does not occur or occurs less than once per week	A	<p>The mandatory evidence is the Wandering/ Verbal/Physical Behaviour Records which will inform on the frequency during the appraisal period.</p> <p>Supporting Evidence:            ACCR or Medical Practitioner notes will inform on related medical and behavioural diagnoses.            Clinical reports can inform on cognition and associated behaviours.            Progress Notes and Care Plans should be congruent with the claim.            Other assessments (Cognition, Behavioural Descriptions List, and modified Behavioural Assessment Form) should be congruent with the claim.</p>
Behaviour occurs at least 2 days in a week	B	
Behaviour occurs at least 6 days in a week	C	
Behaviour occurs twice per day or more, at least six days in a week	D	


To support a B, C or D rating in ACFI 10, the required ACFI assessment is the Cornell Scale of Depression in Dementia (Table 5). It is recommended that users refer to the **ACFI User Guide and Assessment Pack** to support implementation of the tool. The ACFI claim is directly linked to the assessment score and a diagnosis is required for a claim for Rating C or D.

**Table 5: ACFI Question 10 Checklist**

<b>ACFI 10 Symptoms of Depression Checklist</b>	<b>Rating</b>	<b>Where to find the supporting information</b>
Minimal symptoms or symptoms did not occur	A	CSDD assessment score 0-8. Progress Notes, Behavioural Descriptions List and Care Plans should be congruent with the claim.
Symptoms caused mild interference with the person's ability to participate in their regular activities	B	CSDD assessment score 9-13. Progress Notes, Behavioural Descriptions List and Care Plans should be congruent with the claim.
Symptoms caused moderate interference with the person's ability to function and participate in regular activities.	C*	CSDD assessment score 14-18. Progress Notes, Behavioural Descriptions List and Care Plans should be congruent with the claim.
Symptoms of depression caused major interference with the person's ability to function and participate in regular activities	D*	CSDD assessment score 19-38. Progress Notes, Behavioural Descriptions List and Care Plans should be congruent with the claim.
There is a diagnosis or provisional diagnosis of depression completed or reconfirmed in the past 12 months OR Diagnosis or provisional diagnosis of depression being sought and will be made available on request within 3 months of the appraisal date		*Required for a Rating C or D ACCR or Medical Practitioner notes can provide the required diagnoses if they meet the time period criteria.



## Evaluation Process



The evaluation process considers:

- Resident Quality of Life outcomes

Assess if the resident's life is better? In what ways (e.g. happier, healthier)? What might have produced this outcome? This could be determined by revisiting the Quality of Life questionnaire or by interviewing the resident and family.

- Resident Care Outcomes

Have the interventions been successful, for example reduced exiting behaviour?

- Further improvements

What needs re-assessing, what could be implemented in a slightly different way?

## Outcomes

An evaluation of Mr. Grey's care plan has shown:

- *A week later, there has been good progress, Mr. Grey is joining in the music activity, and cooperating with the controlled exercise program. There has been no exiting behaviour noted in the morning.*
- *He is also interacting more with staff and other resident's during the music activity. He appears to be much happier, smiling more, interacting with the environment.*
- *Staff have reported that they are building a better rapport with Mr. Grey during ADLs also, with reduced resistance to staff assistance during ADLs.*
- *He has only attempted to exit the grounds twice during the past week, in the afternoon, when there was no activity in place to distract him.*



## Summary: Steps and Information Flow

Figure 3 shows the phases and steps in the behavioural expression process. It involves:

### **The Identification phase which comprises:**

- Collation of history from documentation and resident/family interviews
- Screening using the Behavioural Descriptions List/Initial Nurse Assessment (limited mood items)
- Completing the Comprehensive Assessment (Behaviour Records, modified Behaviour Assessment Form, modified Cornell Scale for Depression in Dementia, and completing Standardised Care Processes to address any clinical risks)

### **The implementation phase which comprises:**

- Completing referrals as required to fill in assessment gaps or for specialist advice
- Analysing the information to develop strategies based on evidence informed practice
- Setting goals with the resident and family and completing the care plan
- Documenting the evidence steps and completing the ACFI


### **The evaluation phase which comprises:**

- A review of the effectiveness of the interventions on resident outcomes  
Monitoring the progress and effectiveness of the interventions will inform if any changes should be considered
- If a review is required, then at this point the process returns to the identification phase and the associated steps

**The Evidence Based Clinical Assessment Toolkit (EBCAT) is so named because evidence gives us a reason to follow a particular plan toward improved health.**



Figure 3: Behavioural Expressions Summary: Steps and information flow





## Behavioural Expressions Resources

The recommended resources and information on where they are found in the toolkit are listed below (Table 6).

**Table 6: Behavioural Expressions Workbook Resources**

Resource Type	Details	Document
Background Reading	IPA BPSD guidelines- Specialist's guide and Nurses guide DTSC On-line dementia course Module 1	Facility to download/provide access to staff
Screen	Initial Nurse Assessment (e.g. CHA)	Behavioural Expressions (BE) Assessment Pack
Screen	Behavioural Descriptions List	BE Assessment Pack
Screen/Assessment	Modified Behaviour Assessment Form (BAF)	BE Assessment Pack
Mandated Evidence	Behaviour Records	ACFI Assessment Pack (facility to download)
Mandated Evidence	Modified Cornell Scale of Depression in Dementia (CSDD)	ACFI Assessment Pack (facility to download)
Standardised Care Processes (SCP)	Delirium	BE Assessment Pack
Standardised Care Processes (SCP)	Physical Restraint	BE Assessment Pack
Goal setting example	Quality Of Life Questionnaire	Reading Pack
Workbook Exercises	Practice using a Case Study	Behavioural Expressions Workbook Appendix

## Behavioural Expressions References

The recommended resources and references are provided below (Table 7).

**Table 7: References for the Behavioural Expressions Workbook**

Document name	Reference
ACFI Assessment Pack October 2014	Sourced at: <a href="https://www.dss.gov.au/our-responsibilities/ageing-and-aged-care/aged-care-funding/residential-care-subsidy/basic-subsidy-amount-aged-care-funding-instrument/aged-care-funding-instrument-acfi-assessment-pack">https://www.dss.gov.au/our-responsibilities/ageing-and-aged-care/aged-care-funding/residential-care-subsidy/basic-subsidy-amount-aged-care-funding-instrument/aged-care-funding-instrument-acfi-assessment-pack</a>
ACFI User Guide October 2014	Sourced at: <a href="https://www.dss.gov.au/our-responsibilities/ageing-and-aged-care/aged-care-funding/residential-care-subsidy/basic-subsidy-amount-aged-care-funding-instrument/aged-care-funding-instrument-acfi-user-guide">https://www.dss.gov.au/our-responsibilities/ageing-and-aged-care/aged-care-funding/residential-care-subsidy/basic-subsidy-amount-aged-care-funding-instrument/aged-care-funding-instrument-acfi-user-guide</a>
Best care for older people everywhere (BCOPE). The Toolkit (2102).	Department of Health Victoria (2012) Best care for older people everywhere. The toolkit. Sourced at <a href="http://www.health.vic.gov.au/older/toolkit/index.htm">http://www.health.vic.gov.au/older/toolkit/index.htm</a>
Comprehensive Health Assessment (CHA) for Older People in the Health Care System	Authors: La Trobe University ACEBAC Sourced from the Department of Health Victoria The CHA is an example of an initial nurse assessment, it is based on the CHAOP resource.
Comprehensive Health Assessment of the Older Person (CHAOP)	Authors: La Trobe University ACEBAC Sourced from the Department of Health Victoria Resource developed for comprehensive health assessment training for PSRACS (2013).
Dementia Training Study Centres website (DTSC)	10 part on-line dementia education modules. Module 1: Evidence-based tools for BPSD assessment Sourced at: <a href="http://www.dtsc.com.au">http://www.dtsc.com.au</a>
Diagnostic and Statistical Manual of Mental Disorders	American Psychiatric Association: Diagnostic and Statistical Manual of Mental Disorders, Fifth edition. Arlington, VA, American Psychiatric Association, 2013
IPA Complete Guides to Behavioral and Psychological Symptoms of Dementia (BPSD) –Nurses Guide. –Specialists Guide	Sourced at: <a href="http://www.ipa-online.org">www.ipa-online.org</a>
Kozman et al 2006	Authors: Kozman MN, Wattis J, Curran S. Reference: Pharmacological management of behavioural and psychological disturbance in dementia. <i>Hum Psychopharmacol</i> .

Document name	Reference
	2006 Jan; 21(1):1-12.)
NATFRAME Behaviour Assessment Form	Section 7 of the National Framework for Documenting Care in Residential Aged Care Services. Australian Government resource. Sourced at: <a href="http://webarchive.nla.gov.au/gov/20140803082152/http://www.health.gov.au/internet/publications/publishing.nsf/Content/ageing-rescare-natframe.htm~ageing-rescare-natframe01.htm">http://webarchive.nla.gov.au/gov/20140803082152/http://www.health.gov.au/internet/publications/publishing.nsf/Content/ageing-rescare-natframe.htm~ageing-rescare-natframe01.htm</a>
NATFRAME Care profile	Section 11 of the National Framework for Documenting Care in Residential Aged Care Services. Australian Government resource. Sourced at: <a href="https://www.dss.gov.au/our-responsibilities/ageing-and-aged-care/aged-care-funding/residential-care-subsidy/basic-subsidy-amount-aged-care-funding-instrument/suggested-assessment-tools-for-aged-care-funding-instrument-acfi">https://www.dss.gov.au/our-responsibilities/ageing-and-aged-care/aged-care-funding/residential-care-subsidy/basic-subsidy-amount-aged-care-funding-instrument/suggested-assessment-tools-for-aged-care-funding-instrument-acfi</a>
Standardised Care Processes (SCP) Depression & Physical Restraint	Published by the Ageing and Aged Care Branch, Victorian Government, Department of Health, Melbourne, Victoria (2012). Authors: La Trobe University ACEBAC Sourced at: Depression <a href="http://www.health.vic.gov.au/agedcare/downloads/score/depression_scp.pdf">http://www.health.vic.gov.au/agedcare/downloads/score/depression_scp.pdf</a> Physical Restraint <a href="http://www.health.vic.gov.au/agedcare/downloads/score/restraint_scp.pdf">http://www.health.vic.gov.au/agedcare/downloads/score/restraint_scp.pdf</a>

## Behavioural Expressions Workbook Exercises

Now that you have worked through the workbook, you are ready to complete a case study. You may choose to do this by yourself or as a group discussion.

*Facility to insert a case study relevant to ACFI 7 or 8 or 9 or 10*

What clinical history is relevant?

What personal history is relevant?

Start to link the evidence together from your case study

Diagnoses

.....

Body structure/function that is affected

.....

What activities have been restricted

.....

Would you recommend a referral? If yes,

What advice or assessment is missing

.....

Who would the referral be sent to

.....

What information would you prepare for the Health Professional?

.....

What information would you expect from the Health Professional?

.....

Develop two goals with the following aspects:

- Has a single issue focus
- Measurable
- Action orientated strategy
- Realistic and achievable
- Can be evaluated

## Behavioural Expressions Appendix

Resident QoL goal .....
Clinical Care goal .....

What documents do you have, and where are they stored (i.e. documentation trail)

Document Name (insert name)	Location	Date Completed
Resident /family Interview		
ACCR		
CMA/ other medical notes		
Clinical Report		
Initial Assessment		
<input type="checkbox"/> Behavioural Descriptions List <input type="checkbox"/> Behaviour Record (Wandering) <input type="checkbox"/> Behaviour Record (Verbal) <input type="checkbox"/> Behaviour Record (Physical) <input type="checkbox"/> Modified BAF <input type="checkbox"/> Modified CSDD		
Care Plan		
Resident Goals		
Other		

Complete the following ACFI 7/8/9 Checklists and note the evidence for that claim.

ACFI 7/8/9 Checklists	
<i>Behaviour must meet the criteria set out on page 44 of the ACFI User Guide</i>	
<input type="checkbox"/>	<p><b>1 Behaviour does not occur or occurs less than once per week</b></p> <p>Wandering <input type="checkbox"/> [W1]            <input type="checkbox"/> [W2]</p> <p>Verbal        <input type="checkbox"/> [V1]            <input type="checkbox"/> [V2]            <input type="checkbox"/> [V3]            <input type="checkbox"/> [V4]</p> <p>Physical      <input type="checkbox"/> [P1]            <input type="checkbox"/> [P2]            <input type="checkbox"/> [P3]</p>
<input type="checkbox"/>	<p><b>2. Behaviour occurs at least 2 days in a week</b></p> <p>Wandering <input type="checkbox"/> [W1]            <input type="checkbox"/> [W2]</p> <p>Verbal        <input type="checkbox"/> [V1]            <input type="checkbox"/> [V2]            <input type="checkbox"/> [V3]            <input type="checkbox"/> [V4]</p> <p>Physical      <input type="checkbox"/> [P1]            <input type="checkbox"/> [P2]            <input type="checkbox"/> [P3]</p>
<input type="checkbox"/>	<p><b>3. Behaviour occurs at least 6 days in a week</b></p> <p>Wandering <input type="checkbox"/> [W1]            <input type="checkbox"/> [W2]</p> <p>Verbal        <input type="checkbox"/> [V1]            <input type="checkbox"/> [V2]            <input type="checkbox"/> [V3]            <input type="checkbox"/> [V4]</p> <p>Physical      <input type="checkbox"/> [P1]            <input type="checkbox"/> [P2]            <input type="checkbox"/> [P3]</p>
<input type="checkbox"/>	<p><b>3. Behaviour occurs twice a day or more, at least 6 days in a week</b></p> <p>Wandering <input type="checkbox"/> [W1]            <input type="checkbox"/> [W2]</p> <p>Verbal        <input type="checkbox"/> [V1]            <input type="checkbox"/> [V2]            <input type="checkbox"/> [V3]            <input type="checkbox"/> [V4]</p> <p>Physical      <input type="checkbox"/> [P1]            <input type="checkbox"/> [P2]            <input type="checkbox"/> [P3]</p>
<p>Your description of the symptoms for the individual:</p>	

Complete the following ACFI 10 Checklist and note the evidence for that claim.

<b>ACFI 10 Depression Checklists</b> <i>Note: excludes behaviour which is covered in ACFI 8 or ACFI 9. Excludes physical illness or disability as recorded in Medical Diagnosis.</i>	
<input type="checkbox"/>	<b>1 Minimal symptoms or symptoms did not occur</b> <input type="checkbox"/> CSD = 0–8
<input type="checkbox"/>	<b>2. Symptoms caused Mild interference with the person’s ability to participate in their regular activities</b> <input type="checkbox"/> CSD = 9–13
<input type="checkbox"/>	<b>3. Symptoms of depression caused Moderate interference with the person’s ability to function and participate in regular activities</b> <input type="checkbox"/> CSD = 14-18 [Requires a diagnosis for a Rating C]
<input type="checkbox"/>	<b>4. Symptoms caused Major interference with the person’s ability to function and participate in regular activities</b> <input type="checkbox"/> CSD = 19-38 [Requires a diagnosis for a Rating D]
<input type="checkbox"/>	<b>5. There is a diagnosis or provisional diagnosis of depression completed or reconfirmed in the past 12 months</b>
<input type="checkbox"/>	<b>6. Diagnosis or provisional diagnosis of depression being sought and will be made available on request within 3 months of the appraisal date</b>